

## INDIGENOUS SPECIES LIST FOR FORREST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
<i>Acacia melanoxylon</i>	Blackwood	8-20x 5-12m	CSL	●	<b>B, B/F, S/G.</b> Spreading form dense foliage. Screen plant. Prefers deep moist soils. Pale creamy flowers July-Oct. <b>18, 30,201 Widespread coloniser</b>
<i>Acacia mucronata ssp. longifolia</i>	Narrow-leaf Wattle	2-6x 2-5m	SL		<b>B/F S/G.</b> Erect/spreading. Open sites, screen plant. Drought resistant. Narrow leaf, cream flower spike. Aug-Oct. <b>45</b>
<i>Acacia myrtifolia</i>	Myrtle Wattle	3-4x 3-4m	SCL		<b>B/F S/G.</b> Erect/spreading. Open sites, screen plant. Drought resistant. Narrow leaf, cream flower spike. Aug-Oct. <b>45</b>
<i>Acacia nanodealbata</i>	Dwarf Silver Wattle	2-6m			<b>B/F</b> Fast grow. Dry-damp sites. Sun-semi shade. Grey fine leaflets. Bright yellow flowers. Jul-Oct. <b>201</b>
<i>Acacia stricta</i>	Hop Wattle	1-4x 2m	CSL	●	<b>B, B/F</b> Slender, sparse shrub. Drought hardy. Quick grower. Yellow flowers May-Oct. <b>45, 161</b>
<i>Acacia verniciflua</i>	Varnish Wattle	1-4x 3-5m	CS	●	Erect sparsely branched. Hardy. Quick grow. Light screen plant. Wet/dry sites. Young foliage shiny, sticky. Mass yellow flower balls. July-Nov. <b>175,201.</b>
<i>Acacia verticillata ssp. verticillata</i>	Prickly Moses	1-5x 3-5m	CSL	●	<b>B. B/F.</b> Streams – damp shaded wet areas. Prickly low shrub to tree. Yellow flowers June-Dec. <b>45, 201, 161, Widespread coloniser</b>
<i>Acrotriche serrulata</i>	Honey-pots	30cmx .5-1m	CSL		<b>H/E.</b> Slow grow dense g/cover. Damp-well drained sites. Spikey pale yellow tubular flower, honey fragrance. May-Oct. edible fruit. <b>16,21,48.</b>
<i>Allocasuarina paludosa</i>	Scrub Sheoak	1-3m	CSL		Woody, dense erect form. Smooth bark. Branchlets fine grey-green lealess. Poorly drained sites. Jan-Dec. <b>8,165.</b>
<i>Amperea xiphoclada var. xiphoclada</i>	Broom Spurge	30-80x 50cm	SL		Wiry shrub. Dry-damp sites. Rigid stems, few leaves, smooth drk green. Flowers Sep-Dec. <b>16.</b>
<i>Amyema pendula</i>	Drooping Mistletoe				Aerial clumped, drooping form, attaches to host plant mainly Eucalypts, Acacias. Large narrow bronze leaves. Cluster tubular orange-red flowers.
<i>Aotus ericoides</i>	Common Aotus	.5-1.5x .5-1.5m	CSL		<b>B/F.</b> Fast grow. bushy heath like. Tolerates wet/dry periods. Clusters small yellow & red pea flowers. Aug-Dec. <b>16,48.</b>
<i>Astroloma humifusum</i>	Cranberry Heath	30cm- 60cm- 1m	SL		<b>H/E.</b> Dense mat . Damp well drained-dry sites. Sun/part-shade. Sml blue-green stiff leaves. Flower tubular scarlet. Sml. Pale green berry.
<i>Austrocynoglossum latifolium</i>	Forest Hound's-tongue	G/cover r			Damp shaded sites. Prickle like hairs. Tubular flower white, pink, mauve. Flowers Oct-Apr. <b>30,31,201.</b>
<i>Austrodanthonia geniculata</i>	Knead Wallaby-grass	20- 45cm	CSL	●	<b>B/F.</b> Loosly tufted perenn. Flower purplish very slender and drooping. Sep-Feb. <b>45.</b>
<i>Austrodanthonia pilosa</i>	Velvet Wallaby-grass	12cm	CSL		<b>B/F.</b> Erect/flattened tufted perenn. Damp shade sites. Leaves hairy and soft. Flower spikelets short purplish. Nov-Feb. <b>45.</b>
<i>Austrostipa spp.</i>	Spear Grass	50- 100cm	CSL	●	<b>B/F.</b> Perenn rough/hairy leaves. Flower open/dense spikelets. Seeds can penetrate skin, eyes, fleece of animals. Coarse tufted perenn.
<i>Azolla filiculoides</i>	Pacific Azolla	water		●	Aquatic-semi aqu. Perenn. herb. Fan shaped. Leaves triangular, red in sun, green in shade.

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
					Roots free hanging. Two lobed leaves.
<i>Banksia marginata</i>	Silver Banksia	1-8x 1-5m	CSL	●	<b>B/F. H/E. S/G.</b> Heathlands, forests. Leaves toothed dark upper light lower. Honey coloured cylindrical flowers Feb-Jul. Screen plant. <b>161</b>
<i>Baumea acuta</i>	Pale Twig-rush	15- 35cm	SL	●	Slender tufted perenn. Damp-wet sites. Stem flat smooth. Leaves flat. Flower spikelets red-brown. Spr-Sum.
<i>Billardiera scandens var. scandens</i>	Common Apple-berry	climber	CSL		<b>H/E.</b> Dainty climber. Hairy foliage, yellow/green bell flowers. June-Feb. Green edible fruit. <b>16,45,48.</b>
<i>Blechnum minus</i>	Soft Water-fern	1.5m	CSL	●	Hardy. Tufted spreading and erect fronds. Once divided, long narrow. Suit damp sites part shade. Fronds to 1.5m. <b>18,30,31.</b>
<i>Blechnum nudum</i>	Fishbone Water-fern	100x 20cm	CSL	●	Tufted, hardy, quick grow. Light green spreading fronds. Older plants have short trunk. Wet & shade sites. <b>18,30,31,201.</b>
<i>Blechnum wattsii</i>	Hard Water-fern	30- 125cm	SL	●	Hardy creeper/ground cover. Fronds once divided erect dark green, short. Wet areas. <b>18,30,31,201.</b>
<i>Boronia nana</i>	Dwarf Boronia	30cm	CSL		Weak sprawling form. Branchlets hairy. Leaflets 3-5, Flowers pink/white Spring.
<i>Bossiaea cordigera</i>	Wiry Bossiaea	20-50 cmx 1-2.5m	SL		<b>B.</b> Low spreading form, tangled stems. Moist well drained-dry sites. Sun-shade. Tiny heart shape leaves, red, yellow pea flowers. Oct-Jan.
<i>Bossiaea prostrata</i>	Creeping Bossiaea	G/cov	SL	●	<b>B.</b> Low spreading form. Suit most sites but favours damp sites. Stems flattened, hairy. Leaves oval. Flowers red, brown, yellow. Sep-Nov.
<i>Brunonia australis</i>	Blue Pincushion	1m	CSL		<b>B/F.</b> Perenn. herb. Damp well drained sites. Leaves oval, hairy. Flower heads dense, blue tubular, yellow tips stamens. Oct-Jan. <b>22.</b>
<i>Burchardia umbellata</i>	Milkmaids	20- 30cm	CSL		Suit most sites. Grass like leaves, white flowers on stems. Keep dry when dormant. <b>48, 132, 55.</b>
<i>Caladenia s.l. spp.</i>	Orchid	8-30cm	CSL		Leaf solitary at ground hairy. Flowers small like fingers of a hand. solitary or numerous on erect stem.
<i>Callitriche spp.</i>	Water Starwort	Water 90cm	CL		Water/land sites. Creeping mat, small light green rounded leaves, tiny green flowers. <b>Wetlands.</b>
<i>Calochlaena dubia</i>	Common Ground-fern	30- 160cm		●	Erect yellow-green fronds, closely spaced, drooping tips. Sheltered well drained damp/dry sites. <b>18,30,31.</b>
<i>Carex appressa</i>	Tall Sedge	.5-1.5x .5-1m	CSL	●	<b>B/F.</b> Robust, suit wet areas. Full sun. Erect sharp stems, dense flower spike Sept-Nov. <b>18, wetlands</b>
<i>Carex fascicularis</i>	Tassel Sedge	1m	CSL	●	Suit wet sites, full sun. Tufted, leafy mid green, erect stem, drooping dense flower spikes. Sept-Jan. <b>18, wetlands</b>
<i>Cassytha glabella</i>	Slender Dodder-laurel	aerial			Twining herb. Attaching to host plant. Leaves triangular yellow-green. Flower small cluster white. Jun-Oct. <b>8,16.</b>
<i>Cassytha pubescens s.s.</i>	Downy Dodder-laurel	aerial			<b>B/F</b> Weak-robust climber. Attaches to host plant by spikes. Young parts hairy, stems yellow-green – red. Leaves triangular. Flower dense spike. <b>181.</b>
<i>Chamaescilla corymbosa var. corymbosa</i>	Blue Stars	20cm	CSL		Small perenn. herb. Few narrow, green/bluish spreading leaves. Damp sites. Flowers bright blue star. Aug-Nov. Container plant hard to grow in garden. <b>16,48.</b>
<i>Chiloglottis valida</i>	Common Bird-orchid	8-10 cm	CSL		Two dk. broad leaves at base of hairless stem. With purple-green flower resembling birds gaping beak. Oct-Feb. <b>Coloniser</b>
<i>Chiloglottis reflexa</i>	Autumn Bird-orchid	8-15 cm	CSL		Two broad wavy leaves at base of hairless stem. With purple-green flower, resembling gaping beak of bird. Dec-May. <b>Coloniser</b>
<i>Chrysocephalum semipapposum</i>	Clustered Everlasting	80cm	CSL		Perenn. herb. Well drained sunny sites. Leaves narrow, grey-green, sticky. Flowers dense cluster yellow-orange flowers. Oct-Mar.
<i>Clematis aristata</i>	Mountain Clematis	6m Climb	CSL		<b>B.</b> Vigorous dense creeper. Suit most conditions. Cream star flowers. Aug-Mar. Feathery fruits. <b>30, 45, 201</b>

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
<i>Comesperma calymega</i>	Blue-spike Milkwort	60cm	SL		Slender form. Dry sandy sites. Leaves sml. narrow. Flowers pea like, blue terminal clusters Oct-Feb.
<i>Comesperma retusum</i>	Mountain Milkwort	30-160cm	CSL		Erect slender form. Damp sites. Sml. oblong leaves. Flowers pea like, cluster pale pink-magenta. Nov-Jan.
<i>Comesperma volubile</i>	Love Creeper	3m Climb	CSL	●	Slender, nearly leafless twiner. Well drained sites. Flowers pea like, bright blue, clusters. Aug-Dec. <b>21.</b>
<i>Coprosma quadrifida</i>	Prickly Currant-bush	2-4x 1-1.5m	CL	●	<b>B.</b> Hardy. Wet-damp-shade sites. Open erect spiny shrub. Cream tubular flowers Aug-Nov. Edible red berries. Nest site. <b>18, 30, 45, 201</b>
<i>Corybas (s.l.) spp.</i>	Helmet Orchid	30CM	CSL		Tiny with solitary leaf heart shape. Flower hooded purple short stem. Winter colony forming. <b>coloniser</b>
<i>Crassula decumbens</i>	Spreading Crassula	15cm	CSL		Annual. Low spreading form Prefer damp sites. Narrow fleshy leaves. Flowers white/cream.
<i>Crassula helmsii</i>	Swamp Crassula		CSL		Spreading aquatic/semi aquat. Perenn. Tolerates salt. Leaves fleshy, narrow. Flowers white, tinge red. Aug-Nov. <b>wetlands.</b>
<i>Crassula sieberiana</i>	Sieber Crassula	10cm g/cov	CL		Herb. Grows on rock ledges, crevices, damp sites. Leaves small narrow, red stems, tiny cream –pink cluster flowers. Sep-Jun. Regularly flowers.
<i>Cyathea australis</i>	Rough Tree-fern	5-10x 30cm	CSL	●	Adaptable Shade, tolerate sun. Moist-dry sites. Rough trunk, finely divided dark green fronds from apex to 3m. <b>18, 30, 201</b>
<i>Cyperus gunnii</i>	Flecked Flat-sedge	150cm	CSL	●	Tall tufted perenn. Wet areas, waterways. Leaves long, narrow. Dense flat flower spikelets. Spr-Sum.
<i>Cyperus lucidus</i>	Leafy Flat-sedge	1-1.5x 1-1.3m	CSL	●	Robust tussock perenn. Wet sites, waterways. Sun/part shade. Leaves long narrow. Flower spikelets flat. Spr-Sum.
<i>Daviesia leptophylla</i>	Narrow-leaf Bitter-pea	2m	CSL		Erect form. Dry open sites. Leaves thick rigid dull yellow-green. Flowers pea, yellow-brown, red centre. Spr-Sum.
<i>Deyeuxia quadriseta</i>	Reed Bent-grass	1m	CSL	●	Open tufted perenn. Leaves flat-slightly inrolled. Flower dense spike form. Sept-May. <b>16.</b>
<i>Dianella tasmanica</i>	Tasman Flax-lily	.6-1.5 x.5-2m	SL		<b>H/E. B/F.</b> Robust spreading. Damp well drained -wetter fertile soils. Thick shiny leaves. Blue flower yellow centre, blue fruits. Spr-Sum. <b>45, 201.</b>
<i>Dichelachne crinita</i>	Long-hair Plume-grass	20cm	CSL	●	Open tufted perenn. Suit dry and moist sites. Leaves flat green-blue. Flower dense plume like. Oct-Dec. <b>21,22,128.</b>
<i>Dichelachne sciurea spp. agg.</i>	Short-hair Plume-grass		CSL	●	Open tufted perenn. Suit dry and damp sites. Leaves flat green-blue. Flower dense plume like Oct-Dec. <b>21,22,128.</b>
<i>Dichondra repens</i>	Kidney-weed	g/cov	SL	●	Creeping herb. Lawn substitute kidney shaped green leaves. Tiny cream/green flowers Sept-Dec. <b>Widespread.</b>
<i>Dicksonia antarctica</i>	Soft Tree-fern	g/cov	SL	●	Tree fern fibrous trunk Crown widely spreading dark green fronds. Moist/wet slopes gullies. <b>EVC 18,30,201</b>
<i>Dillwynia cinerascens s.l.</i>	Grey Parrot-pea	.3-1.5m	CSL		<b>B/F.</b> Low multi stemmed form. Leaves narrow, hairy. Flowers red, yellow clusters. Sep-Nov.
<i>Dillwynia glaberrima</i>	Smooth Parrot-pea	2m	SL		Spread/erect. Wiry, twiggy form. Good drainage, part shade. Sml. pointy leaves. Flowers yellow, red, dense clusters. Aug-Dec. <b>8,48.</b>
<i>Dipodium roseum s.l.</i>	Rosy Hyacinth-orchid	80cm	CSL		Leafless. Singly/ groups. Reddish-black flower stem. Flower spike many pink-rose flowers. Dec-Feb.
<i>Drosera peltata ssp. auriculata</i>	Tall Sundew	50cm	CL		Slender herb. Thin stems. Rosette pale green-yellow leaves with sticky hairs that trap insects. Flowers white/pale pink. Aug-Dec. <b>48.</b>
<i>Drosera peltata ssp. peltata</i>	Pale Sundew	50cm	CSL		Poorly drained sites. Small round leaves with sticky hairs (catches insects). Erect stem white/pink flowers Aug-Jan.
<i>Drosera whittakeri ssp. aberrans</i>	Scented Sundew	G/cov	CL		Open sites. Leaves rosette green, bronze, spoon shape, with sticky hairs. Large scented white

Soils: C = clay S = sandy L = loamy B = Birds B/F = Butterflies  
H/E = Honey eaters S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
					flowers 3cm. Open sunny days. Jul-Oct.
<i>Echinopogon ovatus</i>	Common Hedgehog-grass	1x 0.2m	SLC	●	Rhizome perennial. Moist sites. Stems erect. Leaves long rough, hairy. Flower dense hairy, oval. Oct-Jan. <b>45</b> .
<i>Eleocharis acuta</i>	Common Spike-sedge	10- 60cm	CSL		Shallow aquatic/semi aquat. Sun – part shade. Leafless green stem, spikelets tip stem. <b>125</b> .
<i>Empodisma minus</i>	Spreading Rope-rush	2m	CSL		Leafless climber – Stems green, thin, wiry. Suit damp-wet sites. Flower spikelets Aug-Mar. <b>8,16,17,48</b> .
<i>Epacris impressa</i> <i>var. impressa</i>	Common Heath	1x.6m	SL		<b>H/E. B/F.</b> Floral emblem Victoria. Stiff prickly shrub. Suit well drained sites. Tubular flowers pink/white. May-Nov. <b>161</b>
<i>Epacris lanuginosa</i>	Woolly-style Heath	60cm	SL		<b>H/E.</b> Slender form, suit moist/wet areas. Dense foliage, small leaves. Cluster sml white flowers. Aug-Jan. <b>8</b> .
<i>Epilobium billardierianum</i>	Variable Willow-herb	.2-1m	CSL		Erect perenn. herb. Cool damp sites. Hairy stems. Leaves oval. Flowers White/purplish pink. Nov-Mar. <b>17,53,18</b> .
<i>Epilobium billardierianum</i> ssp. <i>intermedium</i>	Variable Willow-herb	0.2-1m			Erect form, hairy stems. Cool/damp conditions. Sun in wet areas. Flowers purple, pink/white, hairy. Nov-Mar.
<i>Eragrostis brownii</i>	Common Love-grass	40cm	CSL	●	Coarse, tufted perenn. Suit most sites. Leaves flat, slender, bright green. Flower open pyramid shape. Sept-Apr. <b>widespread</b> .
<i>Eucalyptus aff. willisii</i> (South-western Victoria)	West Coast Peppermint	15m	CSL		<b>B.</b> Hardy. Smooth grey-white bark. Aromatic foliage. Leaves glossy, narrow. Flowers cream Sept-Nov.
<i>Eucalyptus aromaphloia</i>	Scentbark	12-15x 7-12m	CSL		<b>B.</b> Strongly branched – tall, thick rough scented bark. Prefer fertile soils. Leaves bluish-green. White flowers. Jan-Apr. <b>45</b> .
<i>Eucalyptus cypellocarpa</i>	Mountain Grey-gum	10-40x 15m	CSL	●	<b>B/F. H/E.</b> Hardy upright tree dense canopy. Stunted dry soils. Flowers Feb-July. <b>201,30,45</b> .
<i>Eucalyptus globulus</i>	Southern Blue-gum	25-40x 10-20m	CL		<b>H/E.</b> Fast growing tall shelter tree. Smooth trunk. Cream flowers June-Nov. Ribbed silvery fruit Honey source. <b>30, 45</b>
<i>Eucalyptus obliqua</i>	Messmate Stringybark	5-30x 6-20m	CL		<b>B/F. H/E.</b> Grey/brown fibrous bark. Shade/shelter tree. For large areas. Quick hardy grower. Sml. Cream flowers Jan-Mar. Wineglass fruit. <b>30, 45, 161, 201</b>
<i>Eucalyptus ovata</i>	Swamp Gum	8-30x 5-20m	CL	●	<b>H/E, S/G.</b> Excellent in poorly drained soils. Grey white trunk. Creamy flowers May-Nov. <b>3,8,17,18,45,165</b> .
<i>Eucalyptus radiata</i> ssp. <i>radiata</i>	Narrow-leaf Peppermint	6-20x 6-15m	CSL		<b>B/F. H/E.</b> Dense spreading crown, narrow aromatic leaves. Grey-brown bark. Shade. Tree. Oil. Flowers Oct-Jan. <b>45</b>
<i>Eucalyptus viminalis</i>	Manna Gum	10-40x 8-15m	CSL		<b>B/F. H/E. S/G.</b> suits forest, streams. Ribbon bark, white trunk. Koala food. Fast grow, flowers often. <b>18, 45, 201</b>
<i>Eucalyptus yarraensis</i>	Yarra Gum	15-20X 10m	CL		<b>B/F. H/E. S/G.</b> Spreading form. Rough bark short trunk, spreading crown. Suit wetter sites. Grey-green glossy foliage. Flowers sum. <b>16</b> .
<i>Exocarpos cupressiformis</i>	Cherry Ballard	10m	CSL		Suit most sites. Pyramid shaped. Dense bight green/yellow-green triangular foliage. Flower spikes greenish-yellow. Sep-May.
<i>Gahnia sieberiana</i>	Red-fruit Saw-sedge	2-3m	CSL	●	<b>B. B/F. F.</b> Hardy tussock. Suit moist sites, will tolerate dryer sites, sun/shade. . Spreading flowering stems. Nest site. <b>8,16,17,198</b> .
<i>Geranium solanderi</i> s.l.	Austral Cranesbill	.5x .6-1.5m			Perenn. Suit most sites. Hairy lobed leaves. Well drained sites. Flowers pink. Oct-Jan.
<i>Geranium</i> spp.	Crane's Bill	<b>15cm</b>	<b>CSL</b>	●	<b>Perenn. Tufted spreading herb. Hardy. Sun/semi shade. Leaves lobed, hairy. Dec-Apr.</b>
<i>Glossodia major</i>	Wax-lip Orchid	10- 35cm	CSL		Suit drier sites. Single hairy leaf at ground. Slender erect stem, white with purple spots, purple-mauve fragrant flowers. Aug-Nov.
<i>Glyceria australis</i>	Australian Sweet-	1m	CSL	●	Spreading semi-aquatic perenn. Leaves flat, rough

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
	grass				edges. Flower spikelets open form. Oct-Mar. <b>8,16,17,48,125. wetlands.</b>
<i>Gnaphalium indutum.</i>	Tiny cudweed	5-8cm	CSL		Grey-cottony hairy form. Dry sites tolerates dampure. Leaves narrow. Flowers tubular cluster purplish.
<i>Gompholobium ecostatum</i>	Dwarf Wedge-pea	10-45 20- 40cm	CSL		B. Spreading form. Sun-part shade. Damp-dry sites. Narrow bluish leaves. Flower pea- red, yellow. Spr.
<i>Gonocarpus humilis</i>	Shade Raspwort	30cm	CSL		Perenn. Herb. Damp well drained sites. Sun/semi shade. Rough leaves-stem. Sml green/red flower Oct-Dec. <b>3,16,23.</b>
<i>Gonocarpus micranthus ssp. micranthus</i>	Creeping Raspwort	5-10cm	CSL		Short lived creeping perenn. Suit damp-wet sites. Leaves oval serrated. Flowers reddish Oct-Feb..
<i>Gonocarpus tetragynus</i>	Common Raspwort	30cm	CSL		Erect perenn. Herb. Rough leaves, stems. Tiny green/red flowers. Sept-Jan. <b>16,23,198.</b>
<i>Goodenia lanata</i>	Trailing Goodenia	g/cov .6-1m			Trailing carpet form. Suit damp sites. Sun/part shade. Glossy flat, hairy leaves. Flower yellow Spr-Sum. on long stem. <b>16,21.</b>
<i>Goodenia ovata</i>	Hop Goodenia	1-2.5x 1-3m	CSL	●	<b>B/F.</b> Hardy fast grow. Suit coastal-damp sites. Blackberry replacement. Leaves oval. Bright yellow tubular flowers. Spr-sum. <b>45, 30, 201</b>
<i>Goodia lotifolia</i>	Golden Tip	.5x 2-3m	CSL		<b>B/F.</b> Woody, vigorous suckering plant. Screen plant. Suit most sites. Clover shape leaves. Flower yellow Spr.
<i>Grammitis billardierei</i>	Common Finger-fern	7-15X 40- 70cm	CL	●	Tufted strap like fronds dark green. Grows on rocks-logs. Sheltered damp sites. <b>18,30,31.</b>
<i>Gratiola peruviana</i>	Austral Brooklime	15cm	CSL		Perenn. Herb. Wet sites, waterways. Soft leaves, erect stem white/pink flowers. Oct-May.
<i>Grevillea infecunda</i>	Anglesea Grevillea	1x1m	SL		Well drained sites. Large scrambling form. Coarse toothy prickly leaves. Flowers red Oct-Dec.
<i>Gynatrix pulchella s.l.</i>	Hemp Bush	2-4x 1.5-3m	SL	●	Gullies/streambanks. Quick grower. Leaves turn yellow Autumn. Fragrant green/white flowers. Aug-Oct. <b>18</b>
<i>Hedycarya angustifolia</i>	Austral Mulberry	3-7x 4m	L		<b>B.</b> Damp sheltered slopes/gullies. Leaves large glossy, serrated, hard point. Weakly aromatic. Flowers cream fragrant. Aug-Dec. Fruit yellow/orange. <b>18,30,201.</b>
<i>Helichrysum scorpioides</i>	Button Everlasting	.3x .2-.3m			<b>B/F.</b> Hardy perenn. Suit most sites, full sun. Grey foliage, large yellow button flowers Spr-Aut. Dies back. <b>16,48.</b>
<i>Hemarthria uncinata var. uncinata</i>	Mat Grass	50cm	CL	●	Creeping, rigid perenn. Moist heavy sites. Tangled, matted form. Leaves narrow, flat, mid green. Narrow flower spike. Dec-Feb. <b>163.</b>
<i>Hibbertia fasciculata var. prostrata</i>	Bundled Guinea-flower	30-60x 30cm	SL		Erect form. Damp – well drained sites. Soft hairy needle leaves. Flowers regularly, clusters yellow flowers. <b>16,48.</b>
<i>Hibbertia procumbens</i>	Spreading Guinea-flower	G/cov 30cm	SL		Mat forming. Leaves narrow, hairy. Flowers, clusters yellow. Oct-Dec. <b>16,48.</b>
<i>Hibbertia riparia</i>	Erect Guinea-flower	1.2m	CSL		Suit most sites. Leaves narrow. Flowers yellow. Sep-Dec. <b>16,48.</b>
<i>Hibbertia stricta s.l.</i>	Upright Guinea-flower	50x 80cm	SL		Rounded form. Suit most sites. Masses bright yellow flowers. <b>16,48.</b>
<i>Histiopteris incisa</i>	Bat's Wing Fern	0.5-2 x1m	CSL	●	Coarse spreading ground fern. Damp sites. Pale green erect fronds, triangular shape like bats wings. <b>18,30,201.</b>
<i>Hydrocotyle hirta</i>	Hairy Pennywort	G/cov	CSL	●	Mat forming herb. Flowers pale yellow late summer. hairy kidney shaped leaves. <b>Widespread.</b>
<i>Hymenanthera dentata s.l.</i>	Tree Violet	G/cov	CSL	●	<b>B.</b> Rigid shrub. Suit most sites. Tiny Scented yellow bell flowers Sept-Nov. <b>18</b>
<i>Isolepis inundata</i>	Swamp Club-sedge	40cm	SL		Perenn. Suit wet sites. Slender tufted leafless, pale green stems. Small flower spikes stem tip, Sept-Feb. <b>wetlands.</b>

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
<i>Isopogon ceratophyllus</i>	Horny Cone-bush	.2-.4x .5-1.2m			Woody, compact form. Excellent drainage. Rigid prickly leaves. Clusters yellow flowers Sept-Nov.
<i>Joycea lepidopoda</i>	Scaly-foot Wallaby-grass	60cm	CS	●	Tufted perenn. Leaves flat, hairy. Spikelets purple (young). Narrow branched chaffy flowerhead. Nov-Mar.
<i>Joycea pallida</i>	Silvertop Wallaby-grass	.3-1.5 x1m	CSL	●	Spreading tufted perenn. Well drained sites, tolerate some wet. Fine green-bluish leaves. Spikelets straw colour, red tip. Oct-Jan.
<i>Juncus bufonius</i>	Toad Rush	1-30cm	CL	●	Tufted, narrow bright green leaves. Flowers along stems. Perm/ seasonally damp areas. Flowers Oct-Jan. <b>Widespread</b>
<i>Juncus flavidus</i>	Gold Rush	35- 120cm	CSL		Tufted perenn. Suit wet sites. Dull yellow-green to blue-green stems. Flower spikes. Sep-Dec. <b>18, Widespread</b>
<i>Juncus gregiflorus</i>	Green Rush	40- 170cm	CSL		Suit wet sites, dams. Densely tufted perenn. Stems bright green. Flower clusters dark. Nov-Jan. <b>18, widespread</b>
<i>Juncus pallidus</i>	Pale Rush	1.5m	CSL	●	<b>B/F</b> dense tufted, erect. Suit damp/wet sites. Leaves short. Chaffy flowers below stem tip. Oct-Jan. <b>18, Widespread..</b>
<i>Juncus planifolius</i>	Broad-leaf Rush	2-60cm	CSL	●	Tufted form. Suit damp-wet sites. Stems erect. Leaves short, flat. Flower clusters many, red-brown. Sep-Feb. <b>18, Widespread</b>
<i>Juncus procerus</i>	Tall Rush	1-2x .6-1.5m	CSL	●	Dense tufted perenn. Suit damp- wet sites. Stems dull/bright green. Flowers large clusters, golden brown. Dec-Jan. <b>18, Widespread</b>
<i>Juncus subsecundus</i>	Finger rush	80cm	CSL	●	Suit wet sites, full sun. Leafless slender plant. Branched flower spikes. <b>18, Widespread</b>
<i>Kennedia prostrata</i>	Running Postman	1-2.5m G/cov	SL	●	<b>H/E. B/F.</b> Fast grow, creeper. Well drained sites, sun/semi shade. Long slender stems. Leaves grey-green, hairy. Red pea flowers Apr-Dec.
<i>Lagenophora stipitata</i>	Common Bottle-daisy	.05- .25cm	CSL		Perenn. Herb. Damp well drained sites. Leaves rosette on lower stem, Stalk hairy, mauve flowers. Spr-Sum.
<i>Lemna minor s.l.</i>	Common Duckweed	water			Aquatic. Dainty individual floating plants, forms dense mat.. Separate leaflets 3mm. Fish like to eat it. <b>wetlands.</b>
<i>Lepidosperma elatius</i>	Tall Sword-sedge	1-2.5x 1-3m	CS	●	<b>B/F.</b> Hardy clumping form. Broad flattened ribbed leaves. Flower spikes dk brown-grey. Damp sites shade/semi shade. <b>30, 45, 201 wetlands.</b>
<i>Lepidosperma filiforme</i>	Common Rapier-sedge	1-2x .6-1.5m	CSL	●	Clumping perenn. Damp sites, full sun-part shade. Dry sites, part shade. Erect stiff leaves purple-brown. Single spike Flower spikelets. Nov-Mar. <b>48. wetlands</b>
<i>Lepidosperma gunnii</i>	Slender Sword-sedge	15- 40cm	CSL	●	Slender tufted perenn. Damp sites. Stems rigid, smooth. Flower spikelets red-brown Spr-Sum. <b>30, 45, 201 wetlands.</b>
<i>Lepidosperma laterale var. laterale</i>	Variable Sword-sedge	80cm- 1m	CSL	●	Tufted perenn. sedge. Erect flat flower stem. Flower brown slender one side. Sept-Feb. <b>181</b>
<i>Lepidosperma laterale var. majus</i>	Variable Sword-sedge	-3m	CSL	●	Robust tufted perenn. large tussock. Broad leafy mid green. Arched flat stem. Flower spikelets cluster. Damp sheltered sites. Sept-Feb. <b>17,181.</b>
<i>Lepidosperma semiteres</i>	Wire Rapier-sedge	.3-1x .5-1.5m	CSL	●	Tufted perenn. Damp sites, full-part sun. Stems rigid. Leaves short. Flower spikelets red-brown. Spr-Sum. <b>17,181.</b>
<i>Leptospermum continentale</i>	Prickly Tea-tree	1-4x 1-2m	CSL	●	<b>B. B/F.</b> Hardy prickly shrub. Moist/dry sites. Screen plant. White flowers. Oct-Mar. <b>45, 161</b>
<i>Leptospermum lanigerum</i>	Wooly tea tree	2-5x 1-3m	CSL		<b>B. B/F.</b> Hardy prickly shrub. Moist/dry sites. Screen plant. White flowers. Oct-Mar. <b>45, 161</b>
<i>Leptospermum myrsinoides</i>	Heath Tea-tree	1-4x 1-4m	CSL	●	Woody. Sun-part shade. Screen plant. Leaves hairy grey-green. Flowers pale pink Oct-Nov. <b>16,48.</b>
<i>Leptostigma reptans</i>	Dwarf Nertera	10cm G/cover	CSL		Perenn. herb. Mat form. Damp, shade sites. Hairy stems. Flowers tiny cream-green. Nov-Jan.
<i>Leucopogon virgatus</i>	Common Beard-heath	50cm	CSL		Woody. Suit well drained sites, semi shade. Leaves stiff, narrow drk. green, hairy. Flowers dense spikes,

Soils: C = clay S = sandy L = loamy B = Birds B/F = Butterflies  
H/E = Honey eaters S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
					sml. Fragrant, white cup shape Jul-Dec.
<i>Lindsaea linearis</i>	Screw Fern	5-25cm	CSL	●	Creeping form. Dark green fronds, large segments, fan shape. <b>8,16,48.</b>
<i>Lobelia pedunculata s.l.</i>	Matted Pratia	G/cov	CSL	●	Mat forming. Damp sites. Smooth -hairy oval leaves. Red-purple fan shape white, blue,mauve. Flowers regularly.
<i>Lomandra filiformis</i>	Wattle Mat-rush	15-30x 15- 20cm	CS	●	<b>B/F.</b> Hardy tufted perenn. Damp well drained – dry shady sites. Long, narrow leaves. Fine branched flower stems, sml yellow-brown flowers. Oct-Nov. <b>16,45,48.</b>
<i>Lomandra longifolia ssp. exilis</i>	Cluster-headed Mat- rush	50- 100cm	CSL		<b>B/F.</b> Dense tussock. Drier sites. Leaves narrow. Flower spikes yellow-cream. Oct- Dec. <b>45</b>
<i>Lomandra longifolia ssp. longifolia</i>	Spiny-headed Mat- rush	30cm- 1m x 60cm- 1m	SL	●	<b>B/F.</b> Hardy, perenn. Sun/semi shade. Moist-dry sites. Erect-arching foliage. Spikes perfumed cream flowers, orange berries. Aug-Dec. <b>16,21,45.</b>
<i>Lomandra multiflora ssp. multiflora</i>	Many-flowered Mat- rush	.2-.5x g/cov	S	●	<b>B/F.</b> Tufted perenn. Well drained sites. Long narrow leaves. Flowers sml. Yellow brown around stem. May-Dec. <b>16,21,45.</b>
<i>Lotus spp.</i>	Trefoil	60cm	CSL		<b>B.F.</b> Erect-spreading perenn. herb. Drier sites. Divided leaves. Pink/white pea flowers. Sept-Nov. Leaflets.
<i>Luzula meridionalis</i>	Woodrush	30cm	CSL		Tufted perenn. Suit dry-damp sites. Broad grass like leaves. Erect flower stem, clusters of brown chaffy flowers. Sept-Dec.
<i>Lycopodium deuterodensum</i>	Bushy Clubmoss	20-90 cm	SL	●	Creeping rhizome. Moist but well drained open sites. Erect, pale green narrow leaves, non woody plant. Spreading form, <b>coloniser.</b>
<i>Lythrum hyssopifolia</i>	Small Loosestrife	50cm	CSL		Herb. Narrow leaves along erect stems. Flowers tubular deep pink-purple Sept-Feb.
<i>Melaleuca squarrosa</i>	Scented Paperbark	3-10x 2-6m	CSL	●	<b>B/F. H/E.</b> Erect dense form – open sites. Tall in gullies. Low nutrient –wet sites. Shelter tree. Leaves dark green. Clusters cream bottle brush flowers. Oct-Jan. <b>17,165.</b>
<i>Mentha australis</i>	River Mint	.3-.8x .3-1.5m	CL	●	Vigorous, aromatic perenn. herb. Spreading stems. Flower clusters sml white-lilac.
<i>Mentha laxiflora</i>	Forest Mint	.3-.75x .3-1m	CL		Aromatic perenn. herb. Spreading. Suit damp shade sites. Flower clusters sml blue-lilac Nov-Apr.
<i>Microlaena stipoides var. stipoides</i>	Weeping Grass	80x 60cm	CSL	●	<b>B/F.</b> Slender, tufted perenn. Graceful arching stems. Leaves narrow, flat. Flower spikes narrow, delicate. Tolerate shade, grazing. Lawn substitute. <b>3,22,23.</b>
<i>Mitrasacme pilosa var. stuartii</i>	Hairy Mitrewort	30x 10- 50cm	CSL		Tufted perenn. herb. Hairy stems, leaves. Flowers white/cream Sept-Dec.
<i>Monotoca elliptica s.l.</i>	Tree Broom-heath	2-5x .6-2.5m	SL		<b>B/F.</b> Well drained . Sun-semi shade. Screen. Dense sml narrow stiff leaves. Flowers sml white tubular clusters Jun-Sept..
<i>Monotoca scoparia</i>	Prickly Broom-heath	.3-3x .3-1.5m	S		Woody, prickly. Well drained sun-semi shade. Slow grower. Flowers cream fragrant clusters. Feb-Sept.
<i>Myoporum viscosum</i>	Sticky Boobiella	.5- 2x2m	CSL	●	<b>B.</b> Salt tolerant. Screen plant, sun/semi shade. Sticky stems, small white star flowers purple spot. Sept-Dec. <b>coast scrub</b>
<i>Myriophyllum caput-medusae</i>	Coarse Water-milfoil	.3-2m	water		Aquatic perenn. rooted herb. Leaves oblong, purplish. Flowers yellow-red. Oct-Mar. <b>wetlands.</b>
<i>Neopaxia australasica</i>	White Purslane	Water/ land			Wet areas mainly tolerates dry. Dense tufted mat forming. Stems pink. Leaves longer growing in water. <b>Wetlands.</b>
<i>Notelaea ligustrina</i>	Privet Mock-olive	2-6 x3m	CL		Woody, hardy, dense screen plant. Evergreen, narrow leaves. Moist sheltered sites. Small flowers, ornamental berries. <b>18, 30</b>
<i>Notodanthonia semiannularis</i>	Wetland Wallaby- grass		CSL		Sml tussock perenn. damp sites or sandy sites. Flower spikelets. <b>Wetlands.</b>

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
<i>Olearia argophylla</i>	Musk Daisy-bush	3-8x 3-5m	CSL		<b>B. B/F.</b> Fast growing. Wind break. Damp/shade sites. Flowers cream daisy clusters. Oct-Dec. <b>18, 30, 201</b>
<i>Olearia erubescens</i>	Moth Daisy-bush	1-2m	CSL		Spreading form. Drier sites. Branches, underside leaves white hairs, leaves stiff. Flowers daisy, white, yellow Sep-Dec.
<i>Olearia lirata</i>	Snowy Daisy-bush	1-5x 1-3m	SL		<b>B/F.</b> Open shrub. Well drained shaded sites. Quick growing. Snow white daisy flowers. Spring. <b>30, 45, 201</b>
<i>Olearia speciosa</i>	Netted Daisy-bush	1m	CSL		Open form. Cool damp sites. Branches underside leaves hairy. Flowers daisy white, yellow/brown Nov-Jan. <b>21.</b>
<i>Ozothamnus ferrugineus</i>	Tree Everlasting	2-6x 1-3m	SL	●	Dense, fast growing. Gnarled trunk. Moist drained sites. Dark grey-green narrow leaves. Dense cream flower heads. Nov-Feb. <b>45</b>
<i>Patersonia fragilis</i>	Short Purple-flag	10-20x 40cm	S	●	<b>B/F.</b> Perenn iris like herb. Dry-damp sites. Tufts stiff dull bluish-green leaves. Blue-purple flowers. Sep-Jan.
<i>Pelargonium australe</i>	Austral Stork's-bill	60cm	CSL		Low perenn. Suit most sites. Leaves lobed, soft, hairy. Flowers pink/white clusters. Spr-Sum.
<i>Persicaria hydropiper</i>	Water Pepper	1m			Slender erect. Wet areas. Leaves narrow bristled. Flower stem drooping. Pale green-white. Nut dark brown. Jan-May.
<i>Persoonia juniperina</i>	Prickly Geebung	.3-1x .6-.8m			<b>H/E.</b> Erect/round form. Well drained sites. Fine prickly leaves. Yellow tubular flowers Dec-Mar.
<i>Pimelea humilis</i>	Common Rice-flower	.5m	CSL		<b>B/F.</b> Suit most sites. Straggly mound form. Can sucker. Slender hairy branchlets. Leaves narrow green-grey green. Flower clusters creamy white. Sep-Jan. <b>48.</b>
<i>Pimelea linifolia</i>	Slender Rice-flower	30-60x 15- 30cm	CSL		<b>B/F.</b> Suit most sites. Straggly mound form. Can sucker. Slender hairy branchlets. Leaves narrow green-grey green. Flower clusters creamy white. Sep-Jan. <b>48.</b>
<i>Pittosporum bicolor</i>	Banyalla	3-8x 3- 4m	SL	●	<b>B. B/F.</b> Columnar habit. Moist to wet shaded areas. Leaves narrow, yellow bell Aug-Dec. Orange fruit. <b>18, 30.</b>
<i>Plantago varia</i>	Variable Plantain	6-20cm	CSL		Perenn. Suit most sites. Rosette leaves narrow, serrated. Flowers Aug-Apr. <b>widespread.</b>
<i>Platylobium obtusangulum</i>	Common Flat-pea	30-50 cmx 1m			Well drained sites. Small triangular leaves. Red & yellow pea flowers Spring. <b>16,48.</b>
<i>Poa ensiformis</i>	Sword Tussock-grass	1m	CSL	●	<b>B/F.</b> Robust perenn. Loose - dense tussocks. Vigorous in moist shade sites. Leaves rigid, long, narrow, flat, dk green. Flower spikelets open form. Oct-Mar. <b>45, widespread.</b>
<i>Poa labillardierei</i>	Common Tussock-grass	50- 130cm	CL	●	<b>B. B/F.</b> Dense tussock slender green/purple leaves, year long green. Suit wet sites. Flowers Oct-Feb. <b>18, 45</b>
<i>Poa morrisii</i>	Velvet Tussock-grass	50- 90cm		●	<b>B/F.</b> Soft dense perenn. Dry sites. Leaves velvety soft, hairy grey-green. Flower spikelets open form. Oct-Jan. <b>45.</b>
<i>Poa sieberiana</i> var. <i>sieberiana</i>	Grey Tussock-grass	30- 40cm	CSL	●	<b>B/F.</b> Robust perenn. Fine grey leaves green-purplish flower heads. Filter traps. Oct-Feb. <b>161, 45.</b>
<i>Poa tenera</i>	Slender Tussock-grass	20- 50cm	CSL	●	<b>B/F.</b> Spreading tufted grass. Bright green soft leaves. Flower spikelets open form. Oct-Jan. <b>18, 45.</b>
<i>Polystichum proliferum</i>	Mother Shield-fern	1-1.5x 1-1.5m	CSL	●	Clumping fern, forms colonies. Fronds arching 1.5cm. Suit gullies/forest slopes. <b>18,30,201.</b>
<i>Pomaderris aspera</i>	Hazel Pomaderris	3-12x 2-4m	SL		<b>B. B/F.</b> Slender leafy shrub. Suit damp fertile soils, shade. Creamy green flowers. Oct-Dec. Nest site. <b>18, 30, 201</b>
<i>Potamogeton ochreatus</i>	Blunt Pondweed	4.5m	Water		Submerged perenn. herb, flattened branches. <b>Wetlands.</b>
<i>Prasophyllum</i> spp.	Leek Orchid	70cm			Damp sites. Leaf circular solitary. Flowers sml. Pink-

Soils:  
C = clay      S = sandy      L = loamy

B = Birds      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest.  
161 = coast headland scrub. 201 = shrubby wet forest.



FORREST - INDIGENOUS SPECIES LIST

Botanical Name	Common name	Height Width	Soil class	Erosion control	Comments
					white fragrant. Oct-Nov.
<i>Prostanthera lasianthos</i> var. <i>lasianthos</i>	Victorian Christmas-bush	2-6x 1.2-4m	CL		<b>B.</b> Compact form, screen plant, aromatic foliage. Cool areas sun/shade. White flowers with purple spot. Oct-Dec. <b>201,18.</b>
<i>Pteridium esculentum</i>	Austral Bracken	50cmx 1.5m	CSL	●	Hardy. Damp sites. Fronds very divided, leathery, dark green. <b>Widespread.</b>
<i>Pterostylis longifolia</i> s.l.	Tall Greenhood	15- 80cm	CSL		Suit damp open areas. Several long narrow leaves. Flower stem 1-20 green hood like gaping flowers.
<i>Pultenaea daphnoides</i>	Large-leaf Bush-pea	1-3x .5-2m	SL		<b>B/F.</b> Erect form, downy stems. Well drained sites. Leaves wedge shape, pointed tip. Pea flowers dense clusters yellow, red marks Sep-Nov. <b>45</b>
<i>Pultenaea gunnii</i> ssp. <i>gunnii</i>	Golden Bush-pea	.5-1x .4-1m	CS		<b>B/F.</b> Sml. open shrub. Full to part sun. Tiny crowded leaves. Flowers range & yellow pea clusters.
<i>Pultenaea juniperina</i> s.l.	Prickly Bush-pea	1-3x .5- 1.5m	SL		<b>B/F.</b> Erect shrub. Narrow prickly leaves. Clusters of pea shaped yellow/red flowers. Oct-Dec. <b>45</b>
<i>Pyrorchis nigricans</i>	Red-beaks	25cm	CSL		Thick fleshy leaf dark green at ground. Flowers 2-10, with large hood white & red striped. Sep-Nov.
<i>Sambucus gaudichaudiana</i>	White Elderberry	1-2x 1-3.5m	SL		Hardy. Large leaves – Dense white flowers, translucent edible fruit. Damp/dry/salty sites. Oct-Feb. <b>18,201.</b>
<i>Schoenus apogon</i>	Common Bog-sedge	55cm	CSL	●	Tufted form. Seasonally damp-wet sites. Stems erect. Leaves short. Flower cluster spikelets. <b>8, Wetlands.</b>
<i>Schoenus lepidosperma</i>	Slender Bog-sedge	10- 60cm	CSL	●	Tufted perenn. Damp sites. Stems slender erect. Flower spikelets narrow red-brown. Spr-Sum. <b>8,Wetlands</b>
<i>Selaginella uliginosa</i>	Swamp Selaginella	5- 25cm	CSL	●	Perenn. colonies. Damp sites. Stems branching, erect, leafy. <b>8 wetlands.</b>
<i>Solanum laciniatum</i>	Large Kangaroo Apple	3-4x 3-4m	CSL		<b>B.</b> Fast grow, fleshy bush. Suit most sites. Dark green leaves, deeply lobed and entire. Purple flowers. Oval fruit orange. Sept-Mar. <b>Widespread coloniser..</b>
<i>Sphaerolobium minus</i>	Eastern Globe-pea	30-50x 30- 60cm			Erect rush like. Leaves narrow . Flowers cluster spikelets. Damp well drained sites. Sml yellow pea flower Sep-Jan.
<i>Sphaerolobium vimineum</i> s.l.	Leafless Globe-pea	30-60x 30- 60cm	CSL		Sprawling/erect . Damp-dry sites. Full sun/part sun. Suit sand mounds Leafless. Long spikes orange-crimson pea flowers Wint-Spr. <b>48.</b>
<i>Sprengelia incarnata</i>	Pink Swamp-heath	2m	CSL	●	<b>B/F.</b> Stiff stems. Damp –wet sites. Leaves oval stiff, pointed. Flowers pink tubular, dense cluster. Spr&Sum. <b>8.</b>
<i>Spyridium parvifolium</i>	Dusty Miller	1-3x 1-2m	SL		<b>B/F.</b> Hardy. Erect open form. Screen plant. Shade-dry-damp-salty sites. Clusters sml white flowers. July-Nov. Edible fruit. <b>45, 201</b>
<i>Stellaria flaccida</i>	Forest Starwort	G/cov	CSL	●	Creeping herb- sml white daisy flowers. Damp/semi shade. Oct-Feb. <b>18,30,201.</b>
<i>Stellaria pungens</i>	Prickly Starwort	G/Cov	CSL		Spreading untidy perenn. Damp well drained sites. Hairy tangled stems. Leaves prickly br. green sml. flowers white star Oct-Dec. <b>23.</b>
<i>Stylidium graminifolium</i> s.l.	Grass Trigger-plant	.2-.6x .2-.3m	CSL		Suits most sites. Long grass leaves slender stem. Pink flower spike. Insects trigger pollination. Aug-Nov. <b>3,16,45.</b>
<i>Tetrarrhena distichophylla</i>	Hairy Rice-grass	60cm			Scrambling perenn. Leaves narrow (fishbone like) along stem. Flower narrow spike. Nov-Mar.
<i>Tetrarrhena juncea</i>	Forest Wire-grass	Climb – 8m			<b>B/F.</b> Long trailing or climber rough stems. Damp/wet/sun areas. Mauve flower spikes Nov-Apr. <b>30,45,201.</b>
<i>Tetradlea ciliata</i>	Pink-bells	50cm	CSL		Sml leaves around slender hairy erect stem. Flowers drooping pink. Jul-Dec. <b>16,48.</b>
<i>Thelymitra flexuosa</i>	Twisted Sun-orchid	60x 30cm	CSL		Single leaf ribbed, dark green. Well drained sites, damp peat. Scented flower pale/dark blue spots. Sept-Jan.
<i>Thelymitra ixioides</i>	Spotted Sun-orchid	30-	CSL		Single leaf ribbed, dark green. Well drained sites,

Soils: C = clay S = sandy L = loamy B = Birds B/F = Butterflies H/E = Honey eaters S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest. 161 = coast headland scrub. 201 = shrubby wet forest.

FORREST - INDIGENOUS SPECIES LIST

<u>Botanical Name</u>	<u>Common name</u>	<u>Height Width</u>	<u>Soil class</u>	<u>Erosion control</u>	<u>Comments</u>
<i>s.l.</i>		60cm			damp peat. Scented 2-20 blue flowers pale/dark blue spots. Sept-Jan. <b>Widespread.</b>
<i>Thelymitra pauciflora s.l.</i>	Slender Sun-orchid	30-90 cm	CSL		Suit most sites. Stem green-red. Leaves grass like grey-green. Flower pale blue/pink/red-mauve 12-15mm. Oct-Nov.
<i>Todea barbara</i>	Austral King-fern	2-5m	CSL		Robust tree fern like plant. Barrel shape trunk. Covered in brown roots. Fronds from trunk apex. dk. Green glossy Suit wet-waterlogged gullies,,watercourses.
<i>Triglochin procerum s.l.</i>	Water Ribbons	2-3m	Water		Robust, emergent aquatic. Flat strap leaves floating and erect. Still to slow flowing water. Tolerates dry periods. Flower spike green. <b>Wetlands.</b>
<i>Triglochin striatum</i>	Streaked Arrowgrass	3-50cm	CSL	●	Tufted semi-aquatic perenn. herb. Semi erect narrow leaves. Erect flower stem Narrow spike of small green flowers. Flowers regularly. .
<i>Utricularia australis</i>	Yellow Bladderwort	water			Aquatic perenn. Chain of finely divided leaves with bladders that trap insects. Flowers yellow, red-brown spots. Nov-Apr. <b>wetlands</b>
<i>Veronica calycina</i>	Hairy Speedwell	5-50 cm	CSL		Perenn. herb. Suit wet damp sites. Leaves rounded serrated. Hairy erect stems. Flowers pale purple. Oct-Dec.
<i>Viola cleistogamoides</i>	Hidden Violet	G/cov	CSL		Perenn. herb. Suit most sites. Leaves tufted, oval. Flowers cream, purple tinge, bearded. Sept-Nov. <b>48</b>
<i>Viola hederacea</i>	Ivy-leaf Violet	G/Cov	CL	●	Forms dense mat – Suit damp/wet/shade sites. Flowers purple & white. <b>Widespread.</b>
<i>Wahlenbergia gracilis s.l.</i>	Sprawling Bluebell	10-80cm	CSL	●	Tufted perenn. Suit most sites. Stems branched, leafy. Narrow leaves serrated. Flower oval shape blue. Sep-Jan.
<i>Wahlenbergia spp.</i>	Bluebell	30-50 cm	CSL		Well drained sites. Erect slender plant. Flowers pale bluebell. Sep-Nov.
<i>Xanthorrhoea australis</i>	Austral Grass-tree	1-3m	SL		<b>B/F,H/E.</b> Slow grow perenn. Well drained sunny sites. Thick woody trunk. Grassy tuft of leaves. Flower spike long, mass tiny cream fragrant flowers, after fire. Jul-Dec. <b>8,16,48.</b>
<i>Xanthosia dissecta s.l.</i>	Cut-leaf Xanthosia	15cm g/cov	CL		Leaf divided 3 lobed leaflets. Flowers sml cluster white & pale brown. Sept-Nov.
<i>Xyris operculata</i>	Tall Yellow-eye	15-60 cm			Perenn. Suit wet sites. Leaves sword shape. Flowers tubular. <b>8.</b>

Soils:  
C = clay      S = sandy      L = loamy

B = Birds                      B/F = Butterflies  
H/E = Honey eaters      S/G = Sugar Gliders.

EVC = Ecological Vegetation Class. 18 = riparian forest. 30 = wet riparian forest. 45 = shrubby foothill forest. 161 = coast headland scrub. 201 = shrubby wet forest.